



Πύργος Αθηνών, Κτίριο Γ', 2<sup>ος</sup> όροφος, Λεωφ. Μεσογείων 2-4, Αθήνα 115 27

**Τηλ.:** 210 7712792 | **Κινητό:** 697 20 999 11 | **E-mail:** cky@orthosurgery.gr

# Arthroscopic SLAP Repair Protocol

The intent of this protocol is to provide the clinician/therapist/patient with a guideline for post-operative care. It is no means intended to be a substitute for one's clinical decision making regarding the progression of a patient's post-operative course based on their physical exam/findings, individual progress, and/or the presence of post-operative Complications

**Progression to the next phase based on Clinical Criteria and/or Time Frames as Appropriate.**

## **Phase I - Immediate Post Surgical Phase (Weeks 1-4):**

Goals:

- Maintain integrity of repair
- Gradually increase passive range of motion
- Diminish pain and inflammation
- Prevent muscular inhibition

Precautions:

- **No active** range of motion (AROM) of Shoulder
- Maintain arm in sling, remove only for exercise
- No lifting of objects
- No shoulder motion behind back
- No excessive stretching or sudden movements
- No supporting of body weight by hands
- Keep incision clean and dry

Criteria for progression to the next phase (II):

- Passive range of motion (PROM) Flexion to at least 125 degrees
- PROM ER in scapular plane to neutral
- PROM IR in scapular plane to at least 45 degrees
- PROM Abduction to at least 90 degrees in the scapular plane

**DAY 1 TO 6:**

- Abduction brace / sling
- Sleep in brace / sling

- Pendulum Exercises (to begin 3-5 days after surgery)
- Passive ROM
- Flexion to 70 degrees
- Finger, wrist, and elbow AROM
- Patient education: posture, joint protection, positioning, hygiene, etc.
- Cryotherapy for pain and inflammation
- Day 1-2: as much as possible
- Day 3-6: post activity, or for pain
- Replace bandages with band-aids, day 4 or 5

#### **DAY 7 TO 28:**

- Continue use of brace / sling
- MD/PA visit day 10 to remove sutures
- Progress passive ROM
- -Flexion to 90 degrees
- Abduction in the scapular plane
- ER in scapular plane to 30 degrees
- IR in scapular plane
- Continue Elbow, wrist, and finger AROM / resisted
- Cryotherapy as needed for pain control and inflammation

#### **Phase II - Protection Phase (Week 4-10):**

##### **Goals:**

- Allow healing of soft tissue
- Do not overstress healing tissue
- Gradually restore full passive ROM (week 4-5)
- Decrease pain and inflammation

##### **Precautions:**

- No lifting
- No supporting of body weight by hands and arms
- No excessive behind the back movements
- No sudden jerking motions

##### **Criteria for progression to the next phase (III):**

- Full PROM

### **WEEK 5-6:**

- Continue use of sling full time until end of week 4
- Between weeks 4 and 6 may use sling for comfort only
- Initiate active assisted range of motion (AAROM) flexion in supine position
- Progressive passive ROM until approximately Full ROM at Week 5-6.
- This ROM should be PAIN FREE
- Gentle Scapular/glenohumeral joint mobilization to regain full passive ROM
- Begin rotator cuff isometrics
- Continue previous exercises in Phase I as needed
- Continue all precautions
- Initiate prone rowing to neutral arm position
- May use heat prior to ROM exercises
- May use pool (aquatherapy) for light ROM exercises
- Ice after exercise

### **WEEK 6-8:**

- Continue AAROM and stretching exercises
- Initiate active ROM exercises
- Shoulder flexion scapular plane
- Shoulder abduction
- Initiate strengthening program
- External rotation (ER)/Internal rotation (IR) with therabands/sport cord/tubing
- ER Sidelying
- Lateral Raises\*
- Full Can in Scapular Plane\* (avoid empty can abduction exercises at all times)
- Prone Rowing
- Prone Horizontal Abduction
- Prone Extension
- Elbow Flexion
- Elbow Extension

### **Phase III - Intermediate phase (week 10-14):**

#### **Goals:**

- Full AROM (week 10-12)
- Maintain Full PROM
- Dynamic Shoulder Stability

- Gradual restoration of shoulder strength, power, and endurance
- Optimize neuromuscular control
- Gradual return to functional activities

Precautions:

- No heavy lifting of objects (no heavier than 5 lbs.)
- No sudden lifting or pushing activities
- No sudden jerking motions

Criteria for progression to the next phase (IV):

- Able to tolerate the progression to low-level functional activities
- Demonstrates return of strength / dynamic shoulder stability
- Re-establish dynamic shoulder stability
- Demonstrates adequate strength and dynamic stability for progression to higher demanding work/sport specific activities.

**WEEK 10:**

- Continue stretching and passive ROM (as needed)
- Dynamic stabilization exercises
- \*Patient must be able to elevate arm without shoulder or scapular hiking before initiating isotonic; If unable, continue glenohumeral joint exercises

**WEEK 14:**

- Continue all exercise listed above
- Progress to fundamental shoulder exercises

**Phase IV - Advanced strengthening phase (week 16-22):**

Goals:

- Maintain full non-painful active ROM
- Advance conditioning exercises for Enhanced functional use of UE
- Improve muscular strength, power, and endurance
- Gradual return to full functional activities

**WEEK 16:**

- Continue ROM and self-capsular stretching for ROM maintenance
- Continue progression of strengthening
- Advance proprioceptive, neuromuscular activities

**WEEK 20:**

- Continue all exercises listed above
- Continue to perform ROM stretching, if motion is not complete
- Initiate Thrower's Ten if appropriate

**Phase V - Return to activity phase (week 20-26):****Goals:**

- Gradual return to strenuous work activities
- Gradual return to recreational activities
- Gradual return to sport activities

**WEEK 23:**

- Continue strengthening and stretching
- Continue stretching, if motion is tight

**WEEK 26:**

- May initiate interval sport program (i.e. golf, etc.), if appropriate